

Coronavirus COVID-19

BC Centre for Disease Control | BC Ministry of Health



Intubation Checklist for Suspect and Confirmed Cases of COVID-19

Safety of the health care team is paramount and during a pandemic, health care worker safety is prioritised over the patient. Effective communication is key.

Drs. K Romano, N Chima, E Watson, J Atherstone, J Lohser on behalf of Departments of Anesthesia and Critical Care

Intubation Goals

1. Minimise Aerosol Exposure:

- RSI (avoid BMV)
- Inflate Cuff Prior to Ventilation
- Clamp ETT Before Circuit Disconnection

2. Minimise Waste:

· Rationalise Kit

Equipment

Intubation Bag:

- ETT 7.5 w/ Stylet
- Syringe & Gel
- Mcgrath + 4 Blade
- Anchorfast or Tie
- ETT Clamp
- Flex-Tube
- In-Line Suction

Drugs:

- Ketamine 200MG
- Rocuronium 200MG
- Phenylephine X 2
- Propofol 1% 100ML
- Norepinepherine Infusion
- Additional drugs available on request

Crash Bag:

- Bougie
- ETT 6.5, 7.0, 8.0
- X Mcgrath Blade
- LMA 3 & 4 & 5
- DL MAC 3 & 4
- Cricothyroidotomy Set
- ACLS Drug Box & Zoll Defibrillator

Anteroom Bag:

Plan B Kit (Assembled during brief from Crash Bag)

Team Brief (Lead by Anesthesia)

- 1. Team Introduction & Assign Roles
- 2. Patient Allergies/Consent/Code Status
- 3. Empty Pocket Check
- 4. Review Intubation Plan & Prepare Kit
 - Plan A/B/C
 - RSI Sequence & Drug Dosing Recommend 1.5 mg/kg Rocuronium, 1-2mg/kg Ketamine
 - **Prepare Intubation Bag & Drugs**
 - Prepare Anteroom Bag from Crash Bag
 - Trauma Surgeon notified if Necessary
 - **Review Communication Prompts**

- 5. Room Check
 - Ambu-Bag /OPA/HEPA Filter
 - Capnography
 - Suction + Yanker
 - > 2 IV Pumps
 - **IV Access**
 - Ventilator + Closed Suction
 - Plastic Drape
- 6. Cardiac Arrest: AIRWAY THEN **COMPRESSION!**

Team Roles:

Intubator/Room Lead **ANESTHESIOLOGIST**

Drug Administration

Anteroom Runner (PPE ON) RT or RN

Second RN (External, Observer/Chart)


Second Anesthesiologist

Intensivist (External)

RN **Airway Assistant**









Coronavirus COVID-19

BC Centre for Disease Control | BC Ministry of Health



Intubation Checklist for Suspect and Confirmed Cases of COVID-19

Before Room Entry

- Collect Intubation Bag & Drugs
- Collect Anteroom Bag
 - Collect Additional Discussed Equipment +/- CVC, Arterial Line
- DON PPE
- Buddy Check

Pre-Intubation

Ready Intubation Equipment

- McGrath & Blade
- · ETT w/ Stylet & Syringe
- Anchorfast
- Suction
- Ventilator + Closed Suction + Capnography
- Ambu-Bag (No PEEP Valve) + HEPA + Mask + OPA
- Clamp
- Ventilator preset on stand-by

Check Patient

- Position
- Monitors
- IV Access
- · Prepare Infusions

Check Patient

- · Review Intubation Plan
- Address Concerns
- Reinforce Key Points to Minimize Exposure:
 - Avoid BMV
 - · Tolerance of Hypoxemia
 - Clamp ETT for Circuit Disconnect

Intubation

PRE-OXYGENATE 3-5 MIN W/ AMBU-BAG

RSI

TIME 60S + DRAPE

PLAN A

FAIL

(Alert if Sats < 80%)

CALL FOR ANTEROOM BAG +/- SECOND INTUBATOR +/- EXTERNAL CRASH BAG PLAN B

PLAN C

Post-Intubation

Success

- Inflate Cuff
- Attach to Closed Suction + HEPA + Capnography + Ventilator Circuit
- Confirm ETC02
- Secure Tube
- · Initiate Sedation Infusion
- · Initiate Ventilation Strategy
- Insert NGT

Before Exit

- · Ensure Patient Stability
- Discard Disposables
- RT Clean McGrath with AHP Wipes (White Top) then hand off to Anteroom for Second Clean
- · DOFF PPE with Observer

End

- Wash Exposed Areas
- Hot Debrief; Critical Points
- Restock and Clean Kit





